

AIR POLLUTION IN THE BRAZILIAN AMAZON AND SEX SECONDARY RATIO

Pâmela Rodrigues, *Federal University of Mato Grosso, Brazil*

Eliane Ignotti, *University of State of Mato Grosso and National School of Public Health, Brazil*

Sandra Hacon, *National School of Public Health, Brazil*

Karla Longo, *National Institute of Spatial Research, Brazil*

Saulo Freitas, *National Institute of Spatial Research, Brazil*

Background and Aim: Air pollution has been related to changes in the sex ratio in industrial areas. The aim of this study is to analyze the correlation between secondary sex ratio, and exposure to fine particulate matter from biomass burning in the Brazilian Amazon.

Methods: Ecological study of the correlation between the indicators: annual hours (AH%) of fine particulate matter (PM_{2.5}) and secondary sex ratio (SSR) in the Brazilian Amazon region in 2004 and 2005. The database for the exposure indicator, i.e. the percentage of annual hours of PM_{2.5} above the thresholds of 25•g/m³, 50•g/m³ e 80•g/m³, representative of the air pollution levels of concentration of were produced by the Weather Forecasting and Climatic Study Center of the Brazilian Space Research Institute (CPTEC/INPE). To calculate the indicator of RSS we used the database of the Information System of Birth (SINASC) for all municipalities in the Brazilian Amazon.

Results: The states of Rondonia and Mato Grosso presented the highest levels of particulate matter during the study period. The Spearman correlation of every cut-off points for AH% was not statistical significant with the RSS. The averages of RSS were 1.08 and 1.07 in 2004 and 2005 respectively that represents 8 to 7% more males than females. Conclusion: The secondary sex ratios were not related to the levels of pollutants from biomass burning in the Brazilian Amazon during 2004 and 2005.

References:

Ignotti E, Valente JG, Longo K, Freitas S, Hacon SS, Artaxo P.(2010) Impact on human health of particulate matter emitted from burnings in the Brazilian Amazon region Rev Saúde Pública; 44:121-30.

Davis D L, Gottlieb MB, Stampnitzky JR.(1998) Reduced Ratio of Male to Female Births in Several Industrial Countries. A Sentinel Health Indicator? JAMA; 279: 1018 - 23.

Lichtenfels AJFC, Gomes JB, Pieri PC, Miraglia SGK, Hallak J, Saldiva PHN.(2007) Increased levels of air pollution and a decrease in the human and mouse male-to-female ratio in São Paulo, Brazil. Fertility and Sterility; 87: 230 – 32.